# Acer rubrum Seasonally Flooded Forest [Provisional]

COMMON NAME Red Maple Seasonally Flooded Forest SYNONYM Red Maple Seasonally Flooded Flat

PHYSIOGNOMIC CLASS Forest (I)

PHYSIOGNOMIC SUBCLASS
PHYSIOGNOMIC GROUP
PHYSIOGNOMIC SUBGROUP
Cold-deciduous forest (I.B.2)
Natural/Semi-natural (I.B.2.N)

FORMATION Seasonally flooded cold-deciduous forest (I.B.2.N.e)

ALLIANCE Acer rubrum Seasonally Flooded Forest Alliance

CLASSIFICATION CONFIDENCE LEVEL 3

USFWS WETLAND SYSTEM Palustrine

RANGE *Globally* No information

#### Great Smoky Mountains National Park

This vegetation was sampled on the Cades Cove quadrangle, but not on the Mount Le Conte quadrangle. It is not expected to be common in the Park. On the Cades Cove quadrangle, this vegetation was found on a disturbed flat along Abrams Creek, on the west end of the Cades Cove Loop Road [NOTE: photointerpreter polygon 10].

#### ENVIRONMENTAL DESCRIPTION

## Globally

No information

#### Great Smoky Mountains National Park

This is a disturbed, seasonally flooded, flat along a small perennial stream. It is bordered by pastures and fields and was probably grazed and/or cleared in the past. It shows evidence of disturbance by feral hogs.

# MOST ABUNDANT SPECIES

## **Globally**

No information

## Great Smoky Mountains National Park

<u>Stratum</u> <u>Species</u>

Tree canopy Acer rubrum, Fraxinus americana

Herbaceous Microstegium vimineum

## CHARACTERISTIC SPECIES

**Globally**No information

## Great Smoky Mountains National Park

No information

## VEGETATION DESCRIPTION

 ${\it Globally}$ 

This is a placeholder for community association(s) to be developed in this alliance.

# Great Smoky Mountains National Park

#### OTHER NOTEWORTHY SPECIES

The open, even-aged canopy is dominated by *Acer rubrum* and *Fraxinus americana*. Other species in the canopy and subcanopy include *Liriodendron tulipifera* and *Pinus rigida*. There is no shrub stratum and the dense herbaceous stratum is dominated by the invasive exotic grass, *Microstegium vimineum*. Other species in the herb stratum include *Boehmeria cylindrica*, *Brachyelytrum erectum*, *Carex species* (*Carex annectens*, *Carex crinita*, *Carex intumescens*, *Carex lurida*, *Carex rosea*, *Carex* 

# NATIONAL VEGETATION CLASSIFICATION - SOUTHEASTERN UNITED STATES Cades Cove and Mount Le Conte quadrangles, Great Smoky Mountains National Park

April 1999
- Vegetation Descriptions

squarrosa), Chasmanthium laxum var. sessiliflorum, Elymus virginicus, Juncus effusus, Ligusticum canadense, Lycopus virginicus, Polygonum erectum, Polygonum sagittatum, Smilax glauca, Thelypteris noveboracensis, Toxicodendron radicans ssp. radicans, and Viola macloskeyi ssp. pallens.

CONSERVATION RANK

#### RANK JUSTIFICATION

The conservation status of this community has not yet been assessed.

DATABASE CODE CEGL006347

COMMENTS *Globally* 

# Great Smoky Mountains National Park

This vegetation should be mapped at the alliance level. The taxonomic identification of *Fraxinus americana* should be checked in the field. If this tree is *Fraxinus pennsylvanica*, the vegetation may be classified in a different alliance.

REFERENCES

None